



caremountmedical.com

Mark S. Lemert, MD
347 East 37th Street
New York, NY 10016
Phone:(212) 981-7225
Fax: (212) 981-7212

Colonoscopy Preparation Instructions

(must read at least 7 days before exam)

You are scheduled for a colonoscopy at the NYU ACC Endoscopy Center on _____. The facility is located at **240 East 38th Street between 2nd and 3rd avenue on the 23rd floor. You must arrive 60 minutes prior to the scheduled time of the procedure.** Shortly after checking in, the staff of the endoscopy center will direct you from the waiting room to the pre-procedure area. You will then be interviewed by a nurse and meet Dr. Lemert, the gastroenterologist who will be performing the procedure, and the anesthesiologist who will be providing the sedation. Your medical history will be reviewed and questions that you might have will be answered. At that time, please remember to inform us of any medication allergies that you may have. The duration of the procedure and the recovery time afterward are each approximately 30 minutes in duration. You should expect to be at the facility for about 2 hours.

You **MUST** prearrange to have an escort pick you up from the procedure. (*It is not adequate to simply have someone wait for you outside the facility-they need to physically escort you out of the office.*). If needed, you can call Partners In Care at 212-609-7706 to arrange for an aide to pick you up (the cost is about \$80). Failure to have an escort will result in cancellation of your procedure and a \$250 fee. **We also require that no patients drive for 12 hours following the procedure. If you are a female and may potentially be pregnant, please schedule the procedure at a time when you are sure that you are not pregnant.**

You should stop taking nonsteroidal anti-inflammatory medications (NSAIDS) including Motrin, Advil, Aleve, ibuprofen, vitamin E, and fish oil 5 days prior to the procedure. Aspirin should be stopped (unless your cardiologist recommends otherwise) 1 day prior to the exam. Find below a partial list of medications that have a blood-thinning effect:

Aspirin, Aspirin-dipyridomole (Aggrenox)
Clopidogrel (Plavix) Coumadin (warfarin - usually stopped for 3-5 days, please check w/your prescribing doctor - see below)
Pradaxa (Dabigatran)
Dipyridomole (Persantine)
Heparin and Lovenox

Ibuprofen (Advil, Motrin, Nuprin), Meloxicam (Mobic), Naproxen (Aleve, Naprosyn)
Ticlopidine (Ticlid)
Rivaroxaban (Xarelto)
Apixaban (Eliquis)
Prasugrel (Effient)
Ticagrelor (Brillinta)

If you are on any of the above medications you must check with the prescribing doctor to see if you can stop them and for how many days before the colonoscopy and please let Dr. Lemert know what they say.

If you have a cardiac stent, you must continue taking aspirin up until and including the day of the colonoscopy. In accordance with NYU protocol, your exam will be cancelled if you discontinue aspirin. Additionally, if you have any underlying cardiac condition, you must discuss the management of your cardiac medications with your cardiologist at least one week before your exam. If you take Coumadin for a **cardiac condition, stroke prevention or vascular disease**, you should consult your cardiologist, neurologist or internist prior to stopping the medication. If you have **diabetes** and take either insulin or oral medications, dosing may need to be adjusted for the day prior and the day of the procedure. Please consult your internist or endocrinologist regarding this issue.

You should be aware that there is an approximate 0.2% to 1% risk for complications associated with colonoscopy. These risks include, but are not limited to: 1) adverse reactions to medications, 2) gastrointestinal bleeding and 3) colonic perforation that may require operative repair, and 4) aspiration. You should also be aware that a colonoscopy is not 100% accurate and that there is a small possibility of a missed diagnosis of cancer or polyps despite having a competently performed colonoscopy. Alternative methods of colon cancer screening include Virtual Colonoscopy, Barium Enema, Cologard, FIT Stool Test and Flexible Sigmoidoscopy. If you would like to discuss these options or prefer to schedule one of these procedures instead of a colonoscopy, please discuss this at your office visit.

Please call my office (212-981-7225) 3 to 5 days prior to the procedure to confirm your appointment. Please check with your insurance carrier prior to the day of the procedure to confirm coverage of the procedure at the NYU ACC Endoscopy Center. Please let us know if your insurance carrier requires a specific billing code for your procedure. Cancellations less than 3 business days in advance or failure to show for the scheduled procedure may result in a \$250 charge. If any questions arise prior to the procedure, please contact us at 212-981-7225.

Our gastroenterologists recommend the **SUPREP** preparation, which may be purchased from a local pharmacy with a prescription provided with the information materials. You may also want to buy 'baby wipes' for your comfort during the preparation.

PREPARATION DAY: During the day before the procedure you may have a Low-Residue Diet for Breakfast and Lunch Only (no later than 1pm). (A low-residue diet can be made up of Milk, Plain or Vanilla Yogurt, White Bread, Pasta, Cottage Cheese, Cream Cheese, Butter/Margarine, Eggs). After 1pm, you should then restrict your diet to clear liquids (**nothing red or purple**), which include the following:

Strained fruit juices without pulp.

Water.

Clear broth or bouillon.

Coffee or Tea (black).

Gatorade, Kool-Aid, carbonated and non-carbonated soft drinks.

Jell-O and ice popsicles.
Pedialyte.

Step 1: At 4-5pm- pour **One** of the 6 ounce bottles of SUPREP liquid into the mixing container. Add cool drinking water to the 16 ounce line on the container and mix. Drink ALL the liquid in the container. You MUST drink TWO more 16 ounce containers of WATER over the next 1 hour. (Fill the cup up 2 more times with plain water and drink it.). You can drink even more clear liquids if you want and we recommend that you do (clear broth and bouillon are especially good to drink.). If you walk around your home after drinking the prep, it may help the laxative work quicker.

Step 2: At 9-10pm- pour the remaining 6 ounce bottle of SUPREP liquid into the mixing container. Add cool drinking water to the 16 ounce line on the container and mix. Drink ALL the liquid in the container. You MUST drink TWO more 16 ounce containers of WATER over the next 1 hour (fill the cup up 2 more times with plain water and drink them - you have until 3am). You can continue to drink more clear liquids up until 3am.

After starting the bowel preparation, you may have water and clear liquids up to 3 am the morning of the procedure. You may not have anything to drink within 4 hours of the scheduled time of the procedure. You may take cardiac or blood pressure medications with a sip of water the morning of the procedure. Please remember that you cannot eat solid food 1 day prior to and on the day of exam.

Plan to arrive at the NYU ACC Endoscopy Center, 240 East 38th Street, 23rd floor 60 minutes prior to the scheduled time of the procedure. A nurse from the center will contact you 1-2 d prior to the scheduled procedure date to answer any questions, confirm procedure times, and obtain a health history to expedite the admission process. Please feel free to contact the NYU ACC Endoscopy Center at 646-501-9100 with any questions or concerns. **We recommend that you do not wear any jewelry or bring other valuable items to the procedure.** We hope that this process goes as smoothly as possible for you. If there are any questions during any point of the process, feel free to contact us for assistance. Please call the office if you develop a respiratory infection or a cough prior to your procedure as you may need to reschedule the exam. If you call on an evening or a weekend, please make sure the service puts you in contact with the covering physician.